

Expression of CD133 as a cancer stem cell marker in invasive gastric carcinoma

S. Attia, N. Atwan, M. Arafa, R.A. Shahin

Pathology Department, Faculty of Medicine, Mansoura University, Egypt

Summary

Introduction. Gastric cancer is considered to be the fourth most common malignancy all over the world and the second cause of cancer deaths. Regarding the cancer stem cells (CSCs) theory, they are a small group of tumor cells with unrestricted self-renewal and differentiation abilities that help the tumor formation. There is an interest in the utility of CD133 as a promising marker to detect the tumor stem cell population for a variety of solid malignancies including gastric cancer. Tumors that express stem cell marker such as CD133 was found to be more aggressive tumors with poor prognosis and high liability for recurrence. This study aimed to evaluate the immunohistochemical expression of CD133 in invasive gastric carcinoma and study the relation between CD133 immunohistochemical expression and different clinicopathological parameters.

Material and methods. Seventy seven cases of gastric carcinoma were collected from the surgical pathology unit at Gastroenterology Center, Mansoura University, Egypt. CD133 expression in tumor tissue was evaluated by immunohistochemistry.

Results. CD133 expression positively correlated with tumor metastasis and recurrence. Multivariate analysis revealed CD133 positivity to be an independent prognostic factor for tumor recurrence (P value = 0.03).

Conclusion. CD133 is a good marker that can predict tumor recurrence and metastasis in gastric carcinoma. Even though, studies regarding CSCs are still in their initial stages especially those related to CD133 in gastric cancer.

Key words

CD133 • Stem cells • Gastric cancer • Immunohistochemistry

Introduction

Gastric cancer is one of the most aggressive human malignancies. It is considered by the WHO as the fourth most common cancer worldwide and the second leading cause of cancer deaths ¹. According to The Egyptian National Cancer Institute, it is the fourteenth most common cancer representing 1.8% of cases in both sexes ². Tumor recurrence and metastasis are the main causes of death in gastric cancer patients, regardless of surgical intervention and post-operative adjuvant therapy ³. The overall 5-year survival rate is less than 20% as most cases are usually diagnosed late and are incompatible for curative surgery. Therefore, there is a critical need to find more effective methods to eliminate the cancer cells ⁴.

The CSCs correspond to a subpopulation of cells within the tumor defined by self-renewal, asymmetrical division, and differentiation properties, giving rise to the more or less differentiated cells composing the tumor mass ⁵. CSCs were first known in the early half of the 2000s in solid cancer. A treatment targeting CSCs has become a challenge for radical cure of cancer. Therefore, it is important to identify and isolate CSCs using specific markers known as CSCs markers ⁶. Different markers have been found to be expressed on the surface of CSCs. CD133 is one of these markers that has retained much attention and importance ⁷. CD133 receptors on gastrointestinal cancer cells was firstly recognized by Smith et al. ⁸. It was found that an antibody-drug conjugate against this receptor cause delayed growth response in gastric cancer cell ⁹.

How to cite this article: Attia S, Atwan N, Arafa M, et al. *Expression of CD133 as a cancer stem cell marker in invasive gastric carcinoma*. Pathologica 2019;111:18-23. <https://doi.org/10.32074/1591-951X-51-18>

Correspondence: Samar Attia, Department of Pathology, Faculty of Medicine, Mansoura University, 35516 Mansoura, Egypt - E-mail: drsamarattia8@gmail.com

Material and methods

TISSUE SPECIMENS

This retrospective study was performed using gastrectomy specimens from seventy seven cases of gastric carcinomas that were collected during the period from January 2012 until September 2015 from the archive of the pathology department. None of the patients had received preoperative adjuvant therapy, chemotherapy or radiotherapy. The protocol was approved by the Ethical Committee of Mansoura University, and performed in accordance with the principles of the Declaration of Helsinki. Written informed consents were obtained.

HISTOPATHOLOGY

The paraffin blocks were re-cut in 5- μ m-thick sections and stained with hematoxylin and eosin stain and were independently reviewed by two pathologists for confirmation of the diagnosis.

IMMUNOHISTOCHEMISTRY

For immunohistochemical staining for CD133, 5- μ m-thick tissue sections were cut and mounted on coated slides. The sections were deparaffinized in xylene then rehydrated in a series of decreasing concentration of ethanol. After that, heat-induced antigen retrieval was done using pressure cooker and EDTA buffer (PH 9) and immersed in peroxidase-blocking solution (3% hydrogen peroxide) for 15-20 minutes to inhibit endogenous peroxidase activity, the slides were washed in phosphate buffer saline (PBS). The slides were – then – incubated with the Anti-CD133, mouse monoclonal antibody (1:40, clone AC133, Miltenyi Biotech, Bergisch Gladbach, Germany) that was used as primary antibody for CD133 detection for 1 hour at room temperature, followed by incubation with the secondary antibody (poly horseradish peroxidase conjugate for mouse/rabbit) for 15 minutes at room temperature. Diaminobenzedene (DAB) was the chromogen used to visualize the stain. The slides were then counterstained with hematoxylin, dehydrated, coverslipped, and finally mounted with DPX¹⁰. Both normal gastric mucosa and normal kidney tissue were used as replace internal by positive^{11 12}. Negative control was done by substituting the primary antibody with phosphate buffer saline¹³.

ASSESSMENT OF CD133 IMMUNOSTAINING

CD133 show membranous and cytoplasmic pattern of expression¹⁴. We consider the pattern of expression as the more dominant type was seen in the specimen⁶. CD133 results were scored to as the sum of CD133 intensity of staining reaction in tumor cells plus

CD 133 percent of expression in tumor cells. CD133 intensity was graded from zero to three at which (0 for negative expression, 1 for mild intensity, 2 for moderate intensity and 3 for strong intensity) and CD133 percent of expression also was graded from zero to three at which (0 for negative percent, 1 for > 10% of tumor cells, 2 for 10% to < 50% of tumor cells and 3 for \geq 50% of tumor cells)¹⁵. The receiver operating characteristic (ROC) curve was used to set the cut-off point (Fig. 1)^{16 17}. ROC curves facilitate the detection of the threshold value that corresponds to maximum sensitivity with minimal loss of specificity and above which a test result should be considered positive for some outcome¹⁶.

STATISTICAL ANALYSIS

Statistical analysis was done by using IBM SPSS, version 20.0. The relation between CD133 expression and the clinicopathological parameters were assed as a univariate analysis. Multivariate analysis was done using binary logistic regression test to detect the predictors of CD133 expression.

Results

EXAMINATION OF THE CLINICOPATHOLOGICAL DATA

Seventy seven cases of gastric carcinomas were involved in this study. The mean age of the studied cases was 54.44 years, the youngest patient at time of diagnosis was 24 year old while the oldest was 81 year old. The studied cases showed male predominance as it included 45 males (58.4%) and 32 females (41.6%). The tumor size of the excised biopsies ranged from 1 cm to 18 cm in its maximum diameter with mean tumor size was (6.05 \pm 3.3 cm). Grade II adenocarcinoma was the most common histology. The intestinal type was also the predominant according to Laurén classification. Fungating tumor growth pattern was the commonest. The tumor depth of invasion in most of the cases was T3 (75.3%), lymph node metastasis were positive in fifty seven cases (74.0%), while ten cases only (13.0%) out of total number of the studied cases showed distant metastasis. Regarding TNM staging most of the cases were present in stage III (39%). Nineteen cases were positive for lymphovascular invasion (24.7%) and seventeen cases were positive for perineural invasion (22.1%), eleven cases (14.3%) showed tumor recurrence (Tab. I).

IMMUNOHISTOCHEMISTRY

ROC curve analysis was used to determine the clinically relevant threshold for CD133 positivity, the significant cut off was < 3.5, 32 cases showed CD133




Fig. 1. **A)** Grade II adenocarcinoma (H&E x 200 original magnification). **B)** Mucoid adenocarcinoma (H&E x 200 original magnification). **C)** Signet ring carcinoma (H&E x 400 original magnification). **D)** CD133, membranous reaction (luminal) with moderate intensity in 10 > 50% of tumor cells, score 4 (x 400 original magnification). **E)** CD133, cytoplasmic reaction with mild intensity in 10 > 50% of tumor cells, score 3 (x 200 original magnification). **F)** CD133, cytoplasmic reaction with strong intensity in > 50% of tumor cells, score 6 (x 400 original magnification). **G)** Grade I adenocarcinoma of the stomach (H&E x 400 original magnification). **H)** Undifferentiated adenocarcinoma of the stomach (H&E x 400 original magnification). **I)** Perineural invasion by tumor cells (H&E x 100 original magnification). **J)** CD133, membranous reaction (luminal) with strong intensity in > 50% of tumor cells, score 6 (x 400 original magnification). **K)** CD133, cytoplasmic reaction with strong intensity in > 50% of tumor cells, score 6 (x 400 original magnification). **L)** CD133, positive in tumor cells (x 100 original magnification).

Tab. I. Clinicopathological features of the studied cases.

Age	N	%
≤ 55	37	48.1
< 55	40	51.9
Sex		
Male	45	58.4
Female	32	41.6
Tumor size		
Median ± SD	6.05 ± 3.3	
Mean (min-max)	5.0 (1.0-18.0)	
Tumor depth of invasion		
T1	3	3.9
T2	14	18.2
T3	58	75.3
T4	2	2.6
Regional LN metastasis		
Negative (N0)	20	26.0
Positive (N1, N2, N3)	57	74.0
Distant metastasis		
M0	67	87.0
M1	10	13.0
P TNM stage		
I	10	13
II	27	35.1
III	30	39
IV	10	13.0
Lymphovascular invasion		
Positive	19	24.7
Negative	58	75.3
Perineural invasion		
Positive	17	22.1
Negative	60	77.9
Tumor recurrence		
Positive	11	22.2
Negative	66	77.8
Histological type and grade		
Grade 1 adenocarcinoma	5	11.1
Grade2 adenocarcinoma	34	44.1
Grade3 adenocarcinoma	15	19.4
Mucoid adenocarcinoma	6	7.79
Signet ring carcinoma	11	14.28
Undifferentiated carcinoma	6	7.79
Lauren classification		
Intestinal	60	77.9
Diffuse	17	22.1
Ming classification		
Fungating	35	45.5
Infiltrating	14	18.2
Malignant ulcer	28	36.4

score (0-3) that consider negative and about 45 cases showed CD133 score (4-6) that consider positive. Regarding CD133 pattern of expression ten cases were negative. Thirty five cases showed membranous pattern and thirty two cases showed cytoplasmic pattern

Tab. II. Immunohistochemical characteristics.

CD 133 score	N	%
0-3	32	41.6
4-6	45	58.4
CD133 pattern		
Negative	10	13
Membranous	35	45.5
cytoplasmic	32	41.6

(Tab. II). Examples of the staining patterns are shown in Figure 1.

Table III describes the relation between CD133 score and some of the clinico-pathological features. There was a significant relationship between CD133 score and distant metastasis as nine out of ten cases with distant metastasis showed high CD133 score (4-6) with p value = 0.03. Also there was a significance between CD133 score and tumor recurrence as ten out of eleven cases with tumor recurrence showed high CD133 score with P value (0.02). Multivariate analysis using Binary Logistic regression test demonstrated that Cases with tumor metastasis have 7.75 time incidence of expression of CD133 ($P = 0.058$), while cases with tumor recurrence have 8.86 time incidence of CD133 expression ($p = 0.04$), CD133 is an independent prognostic factor for tumor recurrence (Tab. IV).

Discussion

CD133 was considered as one of the most prominent CSC marker that was found to be over expressed in hepatocellular carcinoma, gastric cancers, colorectal cancer, pancreatic cancer and other types of cancer. In the current study, CD133 positivity was found in 58.4% of cases. It was found that presence of CD133 positive staining in minor populations of cells may be indicative of cancer stem-like cells which lead to tumor recurrence and metastasis. Our study aimed to study the relation between CD133 immunohistochemical expression and different clinicopathological parameters.

This study showed a significant relation found between CD133 score and tumor distant metastasis and recurrence and this agreed with Zhao et al. study that reviewed 336 gastric carcinoma cases, 46 cases show distant metastasis 38 cases out of them show CD133 over expression¹⁰. It also agreed with the results of a meta-analysis based study including 26 Studies with a total of 4729 involved cases, the results of that study revealed relation between CD133 over expression and tumor metastasis¹⁸. Also, significant relation was found between CD133 score and tumor

Tab. III. The relation between CD133 score and the clinicopathological parameters.

Clinicopathological parameters	CD 133 0-3 (n = 32)		CD133 4-6 (n = 45)		test of significance
	N	%	n	%	
Age					
≤ 55	14	43.8	23	51.1	$\chi^2 = 0.41$ p = 0.52
< 55	18	56.2	22	48.9	
Sex					
Male	22	68.8	23	51.1	$\chi^2 = 2.39$ p = 0.12
Female	10	31.2	22	48.9	
Tumor size					
Median (Min-Max)	5.0(1.0-18.0)		6.0(1.0-18.0)		Z = 0.037 P = 0.97
Tumor depth of invasion					
T1	2	6.2	1	2.2	MC P = 0.39
T2	4	12.5	10	22.2	
T3	26	81.2	32	17.1	
T4	0	0.0	2	4.4	
Regional LN metastasis					
Positive NO	8	25.0	12	26.7	$\chi^2 = 0.03$ P = 0.87
Negative N1,N2,N3	24	75.0	33	73.3	
Histological type and grade					
Grade 1 adenocarcinoma	0	0.0	5	11.1	MC P = 0.15
Grade2 adenocarcinoma	14	43.8	20	44.4	
Grade3 adenocarcinoma	6	18.8	9	20.0	
Mucoid adenocarcinoma	3	9.4	3	6.7	
Signet ring carcinoma	4	12.5	7	15.6	
Undifferentiated carcinoma	5	15.6	1	2.2	
Lauren classification					
Intestinal	23	71.9	37	82.2	$\chi^2 = 1.16$ p = 0.28
Diffuse	9	28.1	8	17.8	
Ming classification					
Fungating	15	46.9	20	44.4	$\chi^2 = 0.09$ P = 0.95
Infiltrating	6	18.8	8	17.8	
Ulcerative	11	34.4	17	37.8	
Distant metastasis					
M0	31	96.9	36	80.0	$\chi^2 = 4.7$ P = 0.03*
M1	1	3.1	9	20.0	
PTNM					
I-II	15	46.9	22	48.9	χ^2 P = 0.86
III-IV	17	53.1	23	51.1	
Lymphovascular invasion	8	25.0	11	24.4	$\chi^2 = 0.003$ p = 0.96
Perineural invasion	9	28.1	8	17.8	$\chi^2 = 1.16$ p = 0.28
Tumor recurrence					
Negative	31	96.9	35	77.8	$\chi^2 = 5.57$ p = 0.02*
Positive	1	3.1	10	22.2	

χ^2 = Chi-square test * p value significant if < 0.05. MC = Monte Carlo z = Mann Whitney U test.

Tab. IV. Binary logistic regression to detect predictors of CD133 positive cases.

Predictors	Odds ratio	95% CI	P
Tumor recurrence (positive)	8.86	(1.07-73.18)	0.04*
Tumor metastasis (positive)	7.75	(0.93-64.6)	0.058

P: probability * p value significant < 0.05 CI: Confidence interval.

recurrence as ten out of eleven cases with tumor recurrence had score (4-6) with P value = 0.02, it was also

found that CD133 is an independent prognostic factor for tumor recurrence.

This positive relationship between high CD133 expression and tumor aggressiveness suggests that CD133-positive cells have more cancer stem-like cells. These cancer stem-like cells with their self-renewal and differentiation abilities could be responsible for tumor recurrence and metastasis even after traditional treatment¹⁹. It indicates that CD133 may be one of the promising prognostic markers in gastric adenocarcinoma¹⁰.

This study found no significant relation between CD133 score and the patient sex and age. This finding agreed with that of Ishigami et al.; Zhao et al.; Poddar et al.^{20 10 21} and in contrary to that of Nosrati et al. that reported positive correlation between CD133 and age over 65 years with P value > 0.001 that is statistically highly significant¹¹. Moreover, there was insignificant relation between CD133 expression and tumor size, histological type and grade and TNM staging. This finding disagreed with some studies including Ishigami et al., Zhao et al., Nosrati et al. and Poddar et al. that found significant relation between CD133 expression and tumor size, histological type, grade and TNM staging^{20 10 11 21}. This discrepancy in significance may be related to different patient characteristics, different genetics, different races, different environmental factors, and also the application of different scoring methods for CD133 expression. Insignificant relation between lymphovascular invasion and CD133 expression was found in the current study that agreed with that of Nosrati et al.¹¹, but in contrary to that of Ishigami et al.; Yu et al. and Poddar et al. that reported significant relation between CD133 expression and lymphovascular invasion with p value = (0.027), (0.000) and (0.01) respectively that consider significant^{20, 4, 21}. In conclusion CD133 is a good molecular biomarker that can predict tumor recurrence and metastasis in gastric carcinoma. Even though, studies regarding CSCs are still in their initial stages especially those related to CD133 in gastric cancer. Further studies are required to confirm the role of CD133 in tumor recurrence and metastasis and consequently promote finding more appropriate treatment modalities involving targeted gene therapy.

CONFLICT OF INTEREST STATEMENT

None declared.

References

- Edge SB, Compton CC. *The American Joint Committee on Cancer: the 7th edition of the AJCC cancer staging manual and the future of TNM*. Ann Surg Oncol 2010;17:1471-4.
- Ibrahim AS, Khaled HM, Mikhail NN, et al. *Cancer incidence in Egypt: results of the national population-based cancer registry program*. J Cancer Epidemiol 2014;2014:437971.
- Saricanbaz I, Karahacioglu E, Ekinci O, et al. *Prognostic significance of expression of CD133 and Ki-67 in gastric cancer*. Asian Pac J Cancer Prev 2014;15:8215-9.
- Yu JW, Zhang P, Wu JG, et al. *Expressions and clinical significances of CD133 protein and CD133 mRNA in primary lesion of gastric adenocarcinoma*. J Exp Clin Cancer Res 2010;29:141.
- Nguyen PH, Giraud J, Chambonnier L, et al. *Characterization of biomarkers of tumorigenic and chemoresistant cancer stem cells in human gastric carcinoma*. Clin Cancer Res 2017;23:1586-97.
- Hashimoto K, Aoyagi K, Isoe T, et al. *Expression of CD133 in the cytoplasm is associated with cancer progression and poor prognosis in gastric cancer*. Gastric Cancer 2014;17:97-106.
- Mia-Jan K, Jung SY, Kim IY, et al. *CD133 expression is not an independent prognostic factor in stage II and III colorectal cancer but may predict the better outcome in patients with adjuvant therapy*. BMC Cancer 2013;13:166.
- Smith LM, Nesterova A, Ryan MC, et al. *CD133/prominin-1 is a potential therapeutic target for antibody-drug conjugates in hepatocellular and gastric cancers*. Br J Cancer 2008;99:100-9.
- Boegl M, Prinz C. *CD133 expression in different stages of gastric adenocarcinoma*. Br J Cancer 2009;100:1365.
- Zhao P, Li Y, Lu Y. *Aberrant expression of CD133 protein correlates with Ki-67 expression and is a prognostic marker in gastric adenocarcinoma*. BMC Cancer 2010;10:218.
- Nosrati A, Naghshvar F, Khanari S. *Cancer stem cell markers CD44, CD133 in primary gastric adenocarcinoma*. Int J Mol Cell Med 2014;3:279.
- Bussolati B, Bruno S, Grange C, et al. *Isolation of renal progenitor cells from adult human kidney*. Am J Pathol 2005;166:545-55.
- Humphrey PA, Dehner LP, Pfeifer JD. *The Washington manual of surgical pathology*. Philadelphia: Lippincott Williams & Wilkins 2008.
- Chen S, Song X, Chen Z, et al. *CD133 expression and the prognosis of colorectal cancer: a systematic review and meta-analysis*. PLoS One 2013;8:e56380.
- Shamaa, S, Atwan, N, Abdel-Wahab, et al. *Colorectal cancer stem cells: relation to clinico-pathological features and prognosis*. J Am Sci 2016;12:62-8.
- Zlobec I, Steele R, Terracciano L, et al. *Selecting immunohistochemical cut-off scores for novel biomarkers of progression and survival in colorectal cancer*. J Clin Pathol 2007;60:1112-6.
- Ren F, Sheng WQ, Du X. *CD133: a cancer stem cells marker, is used in colorectal cancers*. World J Gastroenterol 2013;19:2603.
- Fu, W. *The clinical value of cancer stem cell markers in gastric cancer*. Clin Oncol 2017;2:1188.
- Vermeulen L, Todaro M, de Sousa Mello F, et al. *Single-cell cloning of colon cancer stem cells reveals a multi-lineage differentiation capacity*. Proc Natl Acad Sci USA 2008;105:13427-32.
- Ishigami S, Ueno S, Arigami T, et al. *Prognostic impact of CD133 expression in gastric carcinoma*. Anticancer Res 2010;30:2453-7.
- Poddar S, D'Cruze L, Halagowder D. *Co-expression of CXCR4 and CD133 in gastric neoplastic tissue and their correlation with clinicopathological factors and prognosis in gastric cancer*. Int J Cancer Ther Oncol 2015;3:3221.